arXiv:1701.03165v1 [math.KT] 11 Jan 2017

A finitely generated group that does not satisfy the Burghelea Conjecture

A. Dranishnikov and M. Hull

Abstract

We construct a finitely generated group that does not satisfy the Burghelea conjecture.

1 Burghelea Conjecture

For an element x in a group G we denote by G_x the centralizer of x.

Computation of the periodic cyclic homology of a group algebra $\mathbb{Q}G$ was performed in [2].

$$PHC_*(\mathbb{Q}G) = \bigoplus_{[x]\in\langle G\rangle^{fin}, n\geq 0} H_{2n+*}(N_x, \mathbb{Q}) \oplus \bigoplus_{\langle G\rangle^{\infty}} T_*(x, \mathbb{Q})$$

where the group $T_*(x, \mathbb{Q})$ fits into the following short exact sequence

$$0 \to \lim_{\leftarrow}{}^1\{H_{*-1+2n}(N_x, \mathbb{Q})\} \to T_*(x, \mathbb{Q}) \to \lim_{\leftarrow}\{H_{*+2n}(N_x, \mathbb{Q})\} \to 0.$$

Here $N_x = G_x/\langle x\rangle$ is the reduced centralizer, $\langle G\rangle$ denotes the set of conjugacy classes on G, $\langle G\rangle^{fin}$ is the set of conjugacy classes of elements of finite order, and $\langle G\rangle^{\infty}$ the set of conjugacy classes of elements of infinite order. The bonding maps in the inverse sequences are the Gysin homomorphisms $S : H_{m+2}(N_x; \mathbb{Q}) \to H_m(N_x; \mathbb{Q})$ corresponding to the fibration $S^1 \to BG_x \to BN_x$.

Conjecture 1.1 (Generalized Burghelea Conjecture)**.** *Let G be a discrete group, then $T_*(x, \mathbb{Q}) = 0$ for all $x \in \langle G\rangle^{\infty}$.*

Burghelea stated the above conjecture for geometrically finite groups [2]. In the same paper he constructed a countable group that does not satisfy the Generalized Burghelea Conjecture.

Theorem 1.2. *There is a finitely generated group G that does not satisfy the Generalized Burghelea Conjecture.*

Our strategy is to show that a torsion-free version of the countable group constructed by Burghelea can be embedded in a finitely generated group in a way that preserves centralizers. This embedding is based on the theory of small cancellation over relatively hyperbolic groups developed by Osin [5].

2 Malnormal Embeddings

For a group G hyperbolic relative to a subgroup H, a subgroup S of G is called *suitable* if S contains two infinite order elements f and g which are not conjugate to any elements of H such that no non-trivial power of f is conjugate to a non-trivial power of g.

The following is a simplification of [5, Theorem 2.4].

Theorem 2.1. *Let G be a torsion-free group hyperbolic relative to a subgroup H, let $t \in G$, and let S be a suitable subgroup of G. Then there exists a group $\overline{G}$ and an epimorphism $\gamma\colon G \to \overline{G}$ such that*

1. *$\gamma|_H$ is injective.*
2. *$\overline{G}$ is hyperbolic relative to H.*
3. *$\gamma(t) \in \gamma(S)$.*
4. *$\gamma(S)$ is a suitable subgroup of $\overline{G}$.*
5. *$\overline{G}$ is torsion-free.*

We inductively apply the previous theorem to construct the desired embedding. This can be extracted from the proof of [5, Theorem 2.6], but since it is not explicitly stated there we include the proof below.

Recall that a subgroup H of a group G is called *malnormal* if for all $x \in G \backslash H$, $x^{-1}Hx \cap H = \{1\}$.

Theorem 2.2. *Let C be a torsion-free countable group. Then there exists a finitely generated group Γ which contains C as a malnormal subgroup.*

Proof. Let $C = \{1 = c_0, c_1, c_2, ...\}$. We inductively define a sequence of quotients as follows: Let $G_0 = C * F$, where $F = F(x, y)$ is the free group on $\{x, y\}$. Then G_0 is hyperbolic relative to C and F is a suitable subgroup of G_0. Suppose now we have constructed a torsion-free group G_i together with an epimorphism $\alpha_i\colon G_0 \twoheadrightarrow G_i$ such that

1. $\alpha_i|_C$ is injective (so we identify C with its image in G_i)
2. G_i is hyperbolic relative to C.
3. $\alpha_i(c_j) \in \alpha_i(F)$ for $1 \leq j \leq i$.
4. $\alpha_i(F)$ is a suitable subgroup of G_i.
5. G_{i+1} is torsion-free.

Given such G_i, we can apply Theorem 2.1 with $S = \alpha_i(F)$ and $t = c_{i+1}$. Let G_{i+1} be the quotient provided by Theorem 2.1, that is $G_{i+1} = \overline{G_i}$. Define $\alpha_{i+1} = \gamma \circ \alpha_i$, where γ is the epimorphism given by Theorem 2.1. The conditions of Theorem 2.1 imply that G_{i+1} satisfies conditions (1)-(5).

Let Γ be the direct limit of the sequence $G_0 \twoheadrightarrow G_1 \twoheadrightarrow G_2...$, that is $\Gamma = G_0/\bigcup \ker(\alpha_i)$. Let $\beta\colon G_0 \to \Gamma$ be the natural quotient map. Note that $\beta|_F$ is surjective by construction. Indeed, G_0 is generated by $C \cup \{x, y\}$ and for each $c_i \in C$, $\alpha_i(c_i) \in \alpha_i(F)$, hence $\beta(c) \in \beta(F)$. Thus Γ is generated by $\{\beta(x), \beta(y)\}$.

Now $\beta|_C$ is injective, so C embeds in Γ; we identify C with its image in Γ. Suppose $x \in \Gamma$ such that $x^{-1}Cx \cap C \neq \{1\}$. Then there exist $g, h \in C \setminus \{1\}$ such that $x^{-1}gxh^{-1} = 1$. Let $\tilde{x} \in G_0$ such that $\beta(\tilde{x}) = x$. Then for some $i \geq 1$, $\tilde{x}^{-1}g\tilde{x}h^{-1} \in \ker \alpha_i$. This means that $\alpha_i(\tilde{x})^{-1}C\alpha_i(\tilde{x}) \cap C \neq \{1\}$. Since G_i is hyperbolic relative to C, C is malnormal in G_i by [5, Lemma 8.3b]. Hence $\alpha_i(\tilde{x}) \in C$, which means that $x = \beta(\tilde{x}) \in C$. Therefore C is malnormal in Γ.

□

3 Modifications of Kan-Thurston Theorem

Baumslag, Dyer, and Heller [1] made the following addendum to the Kan-Thurston [3] theorem.

Theorem 3.1. *For every connected finite simplicial complex L there is a geometrically finite group G and a map $t : K(G, 1) \to L$ that induces an isomorphism of cohomology.*

The proof in [4] of the Kan-Thurson-Baumslag-Dyer-Heller theorem can be extended to produce the following:

Theorem 3.2. *For every connected locally finite simplicial complex L presented as a union of finite subcomplexes $L = \cup L_i$ there is a locally finite complex $K = K(G, 1)$ and a proper map $t : K(G, 1) \to L$ that induces an isomorphism of cohomology and such that $K_i = t^{-1}(L_i) = K(G_i, 1)$ for geometrically finite G_i and the inclusion $K_i \to K_{i+1}$ induces a monomorphism $G_i \to G$.*

Proof of Theorem 1.2. Let L be a locally finite complex representing $K(\mathbb{Z}, 2)$. Let $t : K(G, 1) \to L$ be as in Theorem 3.2. We note that $\lim_{\leftarrow}\{H_{2n}(L; \mathbb{Q}), S\} \neq 0$ where $S : H_{2n+2}(L; \mathbb{Q}) \to H_{2n}(L; \mathbb{Q})$ is the Gysin homomorphism for the canonical S^1-bundle $S^\infty \to \mathbb{C}P^\infty$. Since $G = \cup G_i$ is the union of geometrically finite groups G_i, it is torsion free. Let

$$1 \to \mathbb{Z} = \langle x \rangle \to C \to G \to 1$$

be the central extension extension that corresponds to a generator $a \in H^*(L) = \mathbb{Z}[a]$, $deg(a) = 2$. We apply Theorem 2.2 to obtain a malnormal embedding $C \to \Gamma$ into a finitely generated group. Then $\Gamma_x = C$ and $N_x = \Gamma_x/\langle x \rangle = G$. Then $T(x; \mathbb{Q}) \neq 0$ in the group Γ. □

References

[1] G. Baumslag, E. Dyer and A. Heller, The topology of discrete groups, *J. Pure Appl. Algebra* **16** (1980), 1-47.

[2] D. Burghelea, The cyclic homology of the group rings, *Comment. Math. Helvetici* **60** (1985), 354-365.

[3] D. M. Kan and W. P. Thurston, Every connected space has the homology of a $K(\pi, 1)$, *Topology* **15**, (1976), 253-258.

[4] C.R.F. Mauder, A Short Proof of a Theorem of Kan and Thurston, *Bull. London Math. Soc.* (1981) **13** (4): 325-327.

[5] D. Osin, Small cancellations over relatively hyperbolic groups and embedding theorems, *Ann. of Math.* **172** (2010), no. 1, 1-39.